

# (G,D) - Number of Weak Product and Composition of Graphs

A.Niranjana<sup>1</sup>, K.Palani<sup>2</sup>, S.Kalavathi<sup>3</sup>

<sup>1</sup> PG Mathematics, <sup>2</sup> Department of Mathematics, A.P.C. Mahalaxmi College for Women, Thoothukudi

<sup>3</sup> Research Scholar, Manonmaniam Sundaranar University, Tirunelveli

**Abstract-** (G,D)-number of graphs was introduced by Palani K and Nagarajan A. Let G be a (V,E) graph. A dominating set is a subset D of V such that every vertex in V-D is adjacent to atleast one vertex of D. A (G,D)-set D of G is a subset D of V(G) which is both a dominating and a geodetic set of G. In this paper, we find the (G,D)-number of weak(or kronecker) product and strong product (or composition) of some standard graphs.

## I INTRODUCTION

Graph Theory is an important branch of Mathematics. It has grown rapidly in recent times with a lot of research activities. In 1958, domination was formulized as a theoretical area in graph theory by C. Berge. He referred to the domination number as the coefficient of external stability and denoted as  $\beta(G)$ . In 1962, Ore [7] was the first to use the term 'Domination' number by  $\delta(G)$  and also he introduced the concept of minimal and minimum dominating set of vertices in graph. In 1977, Hedetniemi et al [6] introduced the accepted notation  $\gamma(G)$  to denote the domination number. Let  $G = (V, E)$  be any graph. A dominating set of a graph G is a set D of vertices of G such that every vertex in  $V-D$  is adjacent to atleast one vertex in D. The minimum cardinality among all dominating sets of G is called the domination number of G. It is denoted by  $\gamma(G)$ . The concept of geodominating (or geodetic) set was introduced by Buckley and Harary in [2] and Chartrand, Zhang and Harary in [3, 4, 5]. Let  $u, v \in V(G)$ . A u-v geodesic is a u-v path of length  $d(u, v)$ . A vertex x is said to lie on a u-v geodesic p if x is any vertex on p. A set S of vertices of G is a geodominating (or geodetic) set if every vertex of G lies on an x-y geodesic for some x, y in S. The minimum cardinality of geodominating set is the geodomination (or geodetic) number of G. It is denoted by  $g(G)$ . K. Palani et al [8, 9, 10] introduced the concept (G,D)- set of graphs. A (G,D)- set of graph G is a subset S of vertices of G which is both dominating and geodominating (or geodetic) set of G. A (G,D)- set of G is said to be a minimal (G,D) set of G if no proper subset of S is a (G,D)- set of G. The minimum cardinality of all minimal (G,D)-set of G is called the (G,D)-number of G. It is denoted by  $\gamma_G(G)$ . All graphs considered here are non-trivial, simple and undirected. The order and size of a graph G are denoted by p and q respectively. Let  $G_1(V_1, E_1)$  and  $G_2(V_2, E_2)$  be two graphs. The **weak(or kronecker) product** [1] of  $G_1$  and  $G_2$  denoted by  $G_1 \odot G_2$  has  $V = V_1 \times V_2$  as its vertex set and  $E = \{(u_1, u_2)(v_1, v_2) / (u_1, v_1) \in E_1 \text{ and } (u_2, v_2) \in E_2\}$  as its edge set. The **strong(or composition) product** [1] of  $G_1$  and  $G_2$  is denoted by  $G_1 \boxtimes G_2$  has  $V = V_1 \times V_2$  as its vertex set, and  $u = (u_1, u_2)$  is adjacent with  $v = (v_1,$

$v_2)$  whenever  $u_1$  is adjacent to  $v_1$  (or)  $u_1 = v_1$  and  $u_2$  is adjacent to  $v_2$ . In this paper, we find the (G,D)-number of weak(or kronecker) product and strong product (or composition) of some standard graphs. The following theorems are from [8, 9, 10]

**1.1 Theorem :**  $\gamma_G(K_n) = n$

**1.2 Theorem :**  $\gamma_G(P_n) = 2 + \left\lfloor \frac{n-4}{3} \right\rfloor$

**1.3 Theorem :**  $\gamma_G(C_n) = \left\lfloor \frac{n}{3} \right\rfloor, n \geq 6$ .

**1.4 Theorem:** Any (G,D) set D of a Graph G contains all extreme points of G. In particular, D contains all the end points of G.

## II (G,D) - NUMBER OF WEAK(OR KRONECKER) PRODUCT OF GRAPHS

Here, we find the (G,D)-number of weak product of some standard graphs.

**2.1 Theorem :**  $\gamma_G(P_2 \odot P_n) = 2 \left\lfloor \frac{n+2}{3} \right\rfloor$

**Proof:** Let  $V(P_2) = \{u_1, u_2\}$  and  $V(P_n) = \{v_1, v_2, \dots, v_n\}$ .

Then,  $P_2 \odot P_n$  looks as in figure 2.1

Here,  $P_2 \odot P_n \cong 2P_n$ .

Therefore,  $\gamma_G(P_2 \odot P_n) = \gamma_G(2P_n)$

$= 2\gamma_G(P_n)$

$= 2 \left\lfloor \frac{n+2}{3} \right\rfloor$ .




Figure 2.1

**2.2 Theorem :**  $\gamma_G(P_3 \odot P_n) = n+2$  for  $n \geq 4, n \neq 5$ .

**Proof:** Let  $n \geq 4, n \neq 5$ . Let  $V(P_3) = \{u_1, u_2, u_3\}$  and  $V(P_n) = \{v_1, v_2, \dots, v_n\}$ .

Then,  $P_3 \odot P_n$  looks as in figure 2.2

Obviously,  $S = \{(u_1, v_1), (u_2, v_1), (u_3, v_1), (u_1, v_n), (u_2, v_n), (u_3, v_n)\}$  is a minimum geodetic set of  $P_3 \odot P_n$ . They dominate only the 6 vertices with second coordinate  $v_2$  and  $v_{n-1}$ .




Figure 2.2

To dominate the remaining vertices we need to select either all the vertices in the second row with second co-ordinate not equal to  $v_1, v_2, v_{n-1}$  and  $v_n$  or atleast the same number of vertices from all the three rows to get a minimum (G,D)-set.

Hence,  $S_1 = S \cup \{(u_2, v_3), (u_2, v_4), \dots, (u_2, v_{n-2})\}$  is a minimum (G,D)-set of  $P_3 \odot P_n$ . Therefore,  $\gamma_G(P_3 \odot P_n) = |S_1| = 6+n-4 = n+2$  for  $n \geq 4, n \neq 5$ .

When  $n = 5$ , the graph is as in figure. 2.3




Figure 2.3

In this case  $S_1 = \{(u_1, v_1), (u_2, v_1), (u_3, v_1), (u_1, v_5), (u_2, v_5), (u_3, v_5), (u_2, v_3), (u_2, v_4)\}$  (or)  $S_2 = \{(u_1, v_1), (u_2, v_1), (u_3, v_1), (u_1, v_5), (u_2, v_5), (u_3, v_5), (u_2, v_3), (u_2, v_2)\}$  are the minimum (G,D)-sets.

Therefore,  $\gamma_G(P_3 \odot P_n) = |S_1| = |S_2| = 8$ .

**2.3 Theorem :**

$$\gamma_G(P_4 \odot P_n) = \begin{cases} 4k + 6 & \text{if } n = 4k + 1 \\ 4(k + 1) & \text{if } n = 4k + 2, 4k + 3, 4(k + 1) \end{cases}$$

**Proof:** Let  $V(P_4 \odot P_n) = \{u_1, u_2, \dots, u_n, v_1, v_2, \dots, v_n, w_1, w_2, \dots, w_n, z_1, z_2, \dots, z_n\}$  with  $u_i, v_i, w_i, z_i$  representing the corresponding row elements. Then,  $P_4 \odot P_n$  is as in figure 2.4

Obviously,  $S = \{u_1, v_1, w_1, z_1, u_n, v_n, w_n, z_n\}$  is a geodetic set of  $P_4 \odot P_n$ .

Also, they dominate the 8 vertices in the second and  $(n-1)^{th}$  column elements.

Further, it is observed that selecting last two consecutive vertices among every four consecutive vertices from the second and third row starting with second column element, we could get a dominating set for the remaining elements.

Therefore, to find the vertices to dominate the remaining vertices of  $P_4 \odot P_n$ . We proceed in 4 cases




Figure 2.4




Figure 2.5

**Case (i)** If  $n = 4k + 1$

Here,  $SU\{v_4, v_5, w_4, w_5, v_8, v_9, w_9, \dots, v_{4(k-1)}, v_{4(k-1)+1}, w_{4(k-1)}, w_{4(k-1)+1}, v_{4k}, w_{4k}\}$  is a minimum  $(G,D)$ -set of  $P_4 \odot P_n$  when  $n = 4k+1$ .

$$\begin{aligned} \text{Therefore, } \gamma_G(P_4 \odot P_n) &= |S| = 8 + 4(k-1) + 2 \\ &= 10 + 4(k-1) \\ &= 4k + 6. \end{aligned}$$

**Case (ii)** If  $n = 4k + 2$ .

Here,  $SU\{v_4, v_5, w_4, w_5, v_8, v_9, w_9, \dots, v_{4k}, v_{4k+1}, w_{4k}, w_{4k+1}\}$  is a minimum  $(G,D)$ -set of  $P_4 \odot P_n$  when  $n = 4k+2$ .

$$\text{Therefore, } \gamma_G(P_4 \odot P_n) = |S| = 8 + 4(k) = 4(k + 2).$$

**Case (iii)** If  $n = 4k + 3$ .

Here,  $SU\{v_4, v_5, w_4, w_5, v_8, v_9, w_9, \dots, v_{4k}, v_{4k+1}, w_{4k}, w_{4k+1}\}$  is a minimum  $(G,D)$ -set of  $P_4 \odot P_n$  when  $n = 4k+3$ .

$$\text{Therefore, } \gamma_G(P_4 \odot P_n) = |S| = 8 + 4(k) = 4(k + 2).$$

**Case (iv)** If  $n = 4(k + 1)$ .

Here,  $SU\{v_4, v_5, w_4, w_5, v_8, v_9, w_9, \dots, v_{4k}, v_{4k+1}, w_{4k}, w_{4k+1}\}$  is a minimum  $(G,D)$ -set of  $P_4 \odot P_n$  when  $n = 4(k+1)$ .

$$\text{Therefore, } \gamma_G(P_4 \odot P_n) = |S| = 8 + 4(k) = 4(k + 2).$$

**2.4 Illustration**

Consider  $P_4 \odot P_5$  as in figure 2.5

Here,  $n = 5 = 4k + 1$  where  $k = 1$

$S = \{u_1, v_1, w_1, z_1, u_5, v_5, w_5, z_5\} \cup \{v_4, w_4\}$  is a minimum  $(G,D)$ -set of  $P_4 \odot P_5$ .

$$\text{Therefore, } \gamma_G(P_4 \odot P_5) = |S| = 10 = 4k + 6.$$

**2.5 Illustration**

Consider  $P_4 \odot P_6$  as in figure 2.6

Here,  $n = 6 = 4k + 2$  where  $k = 1$

$S = \{u_1, v_1, w_1, z_1, u_6, v_6, w_6, z_6\} \cup \{v_4, v_5, w_4, w_5\}$  is a minimum  $(G,D)$ -set of  $P_4 \odot P_6$ .

$$\text{Therefore, } \gamma_G(P_4 \odot P_6) = |S| = 12 = 4(k + 2).$$




Figure 2.6

**2.6 Illustration**

Consider  $P_4 \odot P_7$  as in figure 2.7

Here,  $n = 7 = 4k + 3$  where  $k = 1$

$S = \{u_1, v_1, w_1, z_1, u_7, v_7, w_7, z_7\} \cup \{v_4, v_5, w_4, w_5\}$  is a minimum  $(G,D)$ -set of  $P_4 \odot P_7$ .

$$\text{Therefore, } \gamma_G(P_4 \odot P_7) = |S| = 12 = 4(k + 2).$$




Figure 2.7

**2.7 Illustration :**

Consider  $P_4 \odot P_8$  as in figure 2.8




Figure 2.8

Here,  $n = 8 = 4(k + 1)$  where  $k = 1$

$S = \{u_1, v_1, w_1, z_1, u_8, v_8, w_8, z_8\} \cup \{v_4, v_5, w_4, w_5\}$  is a minimum  $(G, D)$ -set of  $P_4 \odot P_8$ .

Therefore,  $\gamma_G(P_4 \odot P_8) = |S| = 12 = 4(k+2)$ .

**2.8 Theorem :**

$$\gamma_G(C_2 \odot C_n) = \begin{cases} 2\lceil n/3 \rceil & \text{if } n \text{ is even} \\ \lceil 2n/3 \rceil & \text{if } n \text{ is odd} \end{cases}; n \geq 6.$$

**Proof:** Let  $V(C_2) = \{u_1, u_2\}$  and  $V(C_n) = \{v_1, v_2, \dots, v_n\}$ .

Then,  $C_2 \odot C_n$  looks as in figure 2.9

$$\text{Here, } C_2 \odot C_n \cong \begin{cases} 2C_n & \text{if } n \text{ is even} \\ C_{2n} & \text{if } n \text{ is odd} \end{cases}$$

Therefore,  $\gamma_G(C_2 \odot C_n)$

$$= \begin{cases} \gamma_G(2C_n) & \text{if } n \text{ is even} \\ \gamma_G(C_{2n}) & \text{if } n \text{ is odd} \end{cases}$$

$$= \begin{cases} 2\gamma_G(C_n) & \text{if } n \text{ is even} \\ \gamma_G(C_{2n}) & \text{if } n \text{ is odd} \end{cases}$$




Figure 2.9

Therefore, by Theorem 1.3,

$$\gamma_G(C_2 \odot C_n) = \begin{cases} 2\lceil n/3 \rceil & \text{if } n \text{ is even} \\ \lceil 2n/3 \rceil & \text{if } n \text{ is odd} \end{cases}$$

**2.9 Remark :**

(1)  $\gamma_G(C_2 \odot C_3) = \gamma_G(C_6) = \lceil 6/3 \rceil = 2$ .

(2)  $\gamma_G(C_2 \odot C_4) = 2 \gamma_G(C_4) = 2 \times 2 = 4$

$$= 2 \times \left\lceil \frac{4}{3} \right\rceil$$

(3) By (1) and (2), it is observed that the above theorem is true for all values of  $n \geq 3$  though  $\gamma_G(C_n) = \lceil n/3 \rceil$  for  $n \geq 6$ .

**2.10 Theorem :**

$$\gamma_G(C_3 \odot C_n) = n; n \geq 3$$

**Proof:** Let  $n \geq 3$ . Let  $V(C_3) = \{u_1, u_2, u_3\}$  and  $V(C_n) = \{v_1, v_2, \dots, v_n\}$ .

Then,  $C_3 \odot C_n$  looks as in figure 2.10 Obviously,  $S = \{(u_1, v_1), (u_1, v_2), (u_1, v_3), (u_1, v_4), \dots, (u_1, v_{n-2}), (u_1, v_{n-1}), (u_1, v_n)\}$  or  $\{(u_2, v_1), (u_2, v_2), (u_2, v_3), (u_2, v_4), \dots, (u_2, v_{n-2}), (u_2, v_{n-1}), (u_2, v_n)\}$  or  $\{(u_3, v_1), (u_3, v_2), (u_3, v_3), (u_3, v_4), \dots, (u_3, v_{n-2}), (u_3, v_{n-1}), (u_3, v_n)\}$  is a minimum geodetic set of  $C_3 \odot C_n$ .

Also, they dominate all the remaining vertices in  $C_3 \odot C_n$ . Thus  $S$  is a minimum  $(G, D)$ -set of  $C_3 \odot C_n$ . Therefore,

$$\gamma_G(C_3 \odot C_n) = n; n \geq 3$$




Figure 2.10

### III (G,D) – NUMBER OF STRONG(OR COMPOSITION) PRODUCT OF GRAPHS

Here, we find the (G,D)-number of strong product of some standard graphs.

**3.1 Proposition :**  $\gamma_G(P_2 \boxtimes P_3) = 2$ .

**Proof:** Let  $V(P_2) = \{u_1, u_2\}$  and  $V(P_3) = \{v_1, v_2, v_3\}$  Then,  $P_2 \boxtimes P_3$  looks as in figure 3.1




Figure 3.1

From figure, clearly  $S = \{(u_1, v_1), (u_1, v_3)\}$  is one of the minimum (G,D)-set of  $P_2 \boxtimes P_3$ .

Therefore,  $\gamma_G(P_2 \boxtimes P_3) = |S| = 2$ .

**3.2 Proposition :**  $\gamma_G(P_2 \boxtimes P_4) = 3$ .

**Proof:** Let  $V(P_2) = \{u_1, u_2\}$  and  $V(P_4) = \{v_1, v_2, v_3, v_4\}$

Then,  $P_2 \boxtimes P_4$  looks as in figure 3.2




Figure 3.2

From figure, clearly  $S = \{(u_1, v_1), (u_1, v_3), (u_1, v_4)\}$  is one of the minimum (G,D)-set of  $P_2 \boxtimes P_4$ .

Therefore,  $\gamma_G(P_2 \boxtimes P_4) = |S| = 3$ .

**3.3 Theorem :**  $\gamma_G(P_2 \boxtimes P_n) = 4$  where  $n > 4$ .

**Proof:** Label the vertices of  $P_2 \boxtimes P_n$  as  $u_1, u_2, \dots, u_n$  and  $v_1, v_2, \dots, v_n$  as in figure 3.3

It is observed that  $P_2 \boxtimes P_n$  contains a bipartite graph  $K_{n,n}$  as its subgraph. Further,  $V(K_{n,n})$  is partitioned into  $V_1, V_2$  with  $G[V_1], G[V_2]$  are path of length  $n$ .

Hence  $V_1, V_2$  contains atleast two non-adjacent vertices. Therefore, a pair of two non-adjacent vertices from  $V_1$  with a pair of two non-adjacent vertices from  $V_2$  forms a geodetic set.




Figure 3.3

Further,  $P_2 \boxtimes P_n$  contains  $K_{n,n}$  implies the above is also a dominating set of  $P_2 \boxtimes P_n$ .

Hence  $S = \{u_i, u_j, v_i, v_j / u_i, u_j \text{ are non-adjacent and } v_i, v_j \text{ are non-adjacent}\}$  is a (G,D)-set of  $P_2 \boxtimes P_n$ . Further,  $S$  is a minimum if  $n > 4$ .

Hence  $\gamma_G(P_2 \boxtimes P_n) = |S| = 4$ .

**3.4 Theorem :**  $\gamma_G(P_3 \boxtimes P_n) = 4$  where  $n > 3$ .

**Proof:** Label the vertices of  $P_3 \boxtimes P_n$  as  $u_1, u_2, \dots, u_n, v_1, v_2, \dots, v_n, w_1, w_2, \dots, w_n$  as in figure 3.4




Figure 3.4

It is observed that  $S = \{u_i, u_j, v_i, v_j / u_i, u_j \text{ are non-adjacent and } v_i, v_j \text{ are non-adjacent}\}$  is one of the  $(G, D)$ -set of  $P_3 \boxtimes P_n$ .

Further,  $S$  is a minimum if  $m > 3$ . Hence  $\gamma_G(P_3 \boxtimes P_n) = |S| = 4$ .

**3.5 Theorem :**  $\gamma_G(P_4 \boxtimes P_n) = 4$  where  $n > 2$ .

**Proof:** Label the vertices of  $P_4 \boxtimes P_n$  as  $u_1, u_2, \dots, u_n, v_1, v_2, \dots, v_n, w_1, w_2, \dots, w_n, z_1, z_2, \dots, z_n$  as in figure 3.5.

It is observed that  $S = \{v_i, v_j, w_i, w_j / v_i, v_j \text{ are non-adjacent and } w_i, w_j \text{ are non-adjacent}\}$  is a  $(G, D)$ -set of  $P_4 \boxtimes P_n$ .

Further,  $S$  is a minimum if  $n > 2$ . Hence  $\gamma_G(P_4 \boxtimes P_n) = |S| = 4$ .




Figure 3.5

**3.6 Theorem :**  $\gamma_G(C_2 \boxtimes C_2) = 4$ .

**Proof:** Let  $V(C_2) = \{u_1, u_2\}$  and  $V(C_2) = \{v_1, v_2\}$

From figure 3.6,  $C_2 \boxtimes C_2 \cong K_4$ .

By Theorem 1.1,  $\gamma_G(C_2 \boxtimes C_2) = \gamma_G(K_4) = 4$ .




Figure 3.6

**3.7 Theorem:**  $\gamma_G(C_2 \boxtimes C_3) = 6$ .

**Proof:** Let  $V(C_2) = \{u_1, u_2\}$  and  $V(C_3) = \{v_1, v_2, v_3\}$

From figure 3.7,  $C_2 \boxtimes C_3 \cong K_6$

By Theorem 1.1,  $\gamma_G(C_2 \boxtimes C_2) = \gamma_G(K_6) = 6$ .




Figure 3.7

**CONCLUSION:**

In general,  $C_m \boxtimes C_n$  is need not be a complete graph.  $(G, D)$ -number of  $C_m \boxtimes C_n$  for any two integers  $m$  and  $n$  could be investigated in a similar manner.

**REFERENCES:**

[1] A note on the connectivity of Kronecker products of graphs-<http://www.science direct.com>>pii.  
 [2] Buckley F, Harary F and Quintas V L, Extremal results on the geodetic number of a graph, Scientia, volume A2 (1988), 17-26.  
 [3] Chartrand G, Harary F and Zhang P, Geodetic sets in graphs, Discussiones Mathematicae Graph theory, 20 (2000), 129-138e.  
 [4] Chartrand G, Harary F and Zhang P, On the Geodetic number of a graph, Networks, Volume 39(1) (2002), 1-6.

[5] Chartrand G, Zhang P and Harary F, Extremal problems in Geodetic graph Theory, Congressus Numerantium 131 (1998), 55-66.

[6] T.W. Haynes, S.T. Hedetniemi and P.J. Slater, Fundamentals Of Domination in graphs, Marcel Decker, Inc., New York 1998.

[7] Ore.O Theory of Graphs, American Mathematical Society Colloquium Publication 38 (American Mathematical Society Providence RI) 1962.

[8] Palani .K and Kalavathi. S, (G,D) – Number of some special graphs, International Journal Of Engineering and Mathematical Sciences January-June 2014, Volume 5, Issue-1, pp.7-15 ISSN(Print) – 2319 – 4537, (Online) – 2319 – 4545.

[9] Palani. K and Nagarajan. A (G,D) – number of graphs, International Journal Of Engineering and Mathematics Research. ISSN 0976 – 5840 Volume 3(2011), pp 285 -299.

[10] Palani. K ,Nagarajan. A and Mahadevan. G, Results connecting domination, geodetic and (G,D)– number of graph, International Journal Of Combinatorial graph theory and applications, Volume 3, No.1, January – June (2010)(pp.51 -59).

LIBRARY